

6. DAFTAR PUSTAKA

- Abiso, E. Satheesh, N. Addisalem, H. (2015). Effect of Storage Methods and Ripening Stage on Postharvest Quality of Tomato (*Lycopersicum esculentum* Mill) CV. Chali. Food Science and Technology. Vol 16(1).
- Al-Haq, M. I., Sugiyama, J., Isobe, S. (2005). Applications of electrolyzed water in agriculture and food industries. Food Science and Technology Research 11: 135–150.
- Arah, I.K. Amaglo, H. Ernest, K, K. Hayford, O. (2015). Preharvest and Postharvest Factors Affecting the Quality and Shelf Life of Harvested Tomatoes. International Journal of Agronomy. Hindawi.
- Aryal, S. (2016). Spread Plate Technique-Principle, Procedure and Uses. Microbiology Info. Nepal.
- Badan Pusat Statistik Provinsi Jawa Tengah. (2016). Statistik Pertanian Hortikultura Provinsi Jawa Tengah 2014-2016. Semarang, Jawa Tengah.
- Beckle, D. M. (2012). Factors Affecting The Postharvest Soluble Solids And Sugar Content of Tomato (*Solanum lycopersicum* L.) Fruit. Elsevier. Department of Plant Sciences. University of California, United States. Vol 63:129–140.
- Critzer, F. Wszelaki, A. (2012). Using Peroxyacetic Acid (PAA) In Fruit and Vegetable Washing and Packing. Department of Food Science. University of Tennessee.
- Cserhalmi, Z. Sass-Kissb. Tóth-Markusb. Lechnera (2006). Study of pulsed electric field treated citrus juices. Innovative Food Science and Emerging Technologies, 7 (2006), pp. 49-54.
- De Cort, S and Godts, F. (2017). Packaging Materials: Polyethylene Terephthalate (PET) For Food Packaging Applications. International Life Sciences Institute. Europe.
- Environmental Protection Agency (EPA). (2011). *Water Treatment Manual : Disinfection*. Environmental Protection Agency. Wexford, Ireland.
- Ferreira, M, D. Carmelo, L, G, P. Polina, C, S. (2013). Evaluation of Impact Damage during Handling of Fresh Market Tomatoes. Horticultural Society; 126:203-205.
- Focus. (2002). Dissolved Oxygen and the Water Quality Standards. Department of Ecology. Washington.
- Gharezi, M. Joshi, N. Elnaz, S. (2012). Effect of Post Harvest Treatment on Stored Cherry Tomatoes. Nutrition and Food Science. University of Agricultural Science, Bengaluru, India. Vol 2:8.

- Hati. S., Surajit. M., P. S. Minz., Shilpa. V., Yogesh. K., B. P. Sigh., Dipika. Y. (2012). Electrolyzed Oxidized Water (EOW): Non-Thermal Approach for Decontamination of Food Borne Microorganisms in Food Industry. Food and Nutrition Science;3: 760-768.
- Hidayanto, E. Rofiq, A. Heri, S. (2010). Aplikasi Portable Brix Meter untuk Pengukuran Indeks Bias. Berkala Fisika. Universitas Diponegoro, Semarang. Vol 4:113-118.
- Hocking, A,D. Pitt, J,I. R,A, Samson, U,Thrane. (2006). Advance in Food Mycology. Springer, Vol 571.
- Hricova D, Stephan R, Zweifel C. (2008). Electrolyzed water and its application in the food industry. Journal of Food Production; 71(9): 1934-1947.
- Huang, C, C. Yang, C, M. Skyi, K, O. Junne, J, C. (2010). Technology on Rducing Post-harvest Losses and Maintaining Quality of Fruits and Vegetables. Taiwan Agricultural Research Institute. Taiwan.
- Inggas, M.A.N., I M. S. Utama, G. Arda. 2013. Pengaruh Emulsi Minyak Nabati Sebagai Bahan Pelapis pada Buah Tomat (*Lycopersicon esculentum* Mill.) terhadap Mutu dan Masa Simpannya. Universitas Udayana.
- Issa-Zacharia, A. Kamitani, Y. Happiness, S, M. Bernadette, K, N. (2010). A review of microbiological safety of fruits andvegetables and the introduction of *electrolyzed water*as an alternative to sodium hypochlorite solution. African Journal of Food Science. Vol 4(13);778-789.
- Kader, A, A. Kitinoja, L. Awad, M, H. Omar, A. Amer, J. Ahmed, E, S. (2012). Role of Agro-industry In Reducing Food Losses In The Middle East and North Africa Region. Food and Agriculture Organization of the United Nations. Cairo.
- Khaleghi, S, S. Ansari, N, A. Mortazavi, S, M, H. (2013). Effect of Calyx Removal and Disinfection On Ripening Rate and Control of Postharvest Decay of Tomato Fruit. South Western journal of Horticulture, Biology and Enviroment. Vol 4 (2); 103-115.
- Koide, S. Takeda, J. I, Shi, J. Shono, H. Atungulu, G. (2009). Disinfection efficacy of slightly acidic electrolyzed water on fresh cut cabbage. Food Control 20: 294–297.
- Koseki S, Yoshida K, Kamitani Y, Itoh K. (2003). Influence of inoculation method, spot inoculation site, and inoculation size on the efficacy of acidic electrolyzed water against pathogens on lettuce. J Food Prot 66(11):2010-2016.
- Koseki, S. Kyoichiro, Y. Seichiro, I. Kazuhiko, I. (2001). Decontamination of Lettuce Using Acidic Electrolyzed water. Journal of Food Protection; 64(5): 652-658.

- Koseki, S. Kyoichiro, Y. Seiichiro, I. Kazuhiko, I. (2004). Efficiency of Acidic *Electrolyzed water* for Microbial Decontamination of Cucumbers and Strawberries. *Journal of Food Protection*; 67(6): 1247-1251.
- Leide, J. Hildebrandt, U. Kerstin, R. Markus, R. Gerd, V. (2007). The Developmental Pattern of Tomato Fruit Wax Accumulation and Its Impact on Cuticular Transpiration Barrier Properties: Effects of a Deficiency in a b-Ketoacyl-Coenzyme A Synthase (LeCER6). *American Society of Plant Biologies*. Vol. 144, pp. 1667–1679.
- Len, S., Hung, Y., Chung, D., Anderson, J.L., Erickson, M.C., dan Morita, K. (2002), Effects of Storage Conditions and pH on Chlorine Loss in Electrolyzed Oxidizing (EO) Water. *Journal of Agricultural and Food Chemistry* Vol 50(1):209-212.
- Mahajan PV, Caleb OJ, Singh Z, Watkins CB, Geyer M. (2014). Postharvest Treatments Of Fresh Produce. *Phil. Trans. R.*
- Mbajiuka, S. Chinedu. Enya. (2014). Isolation of Microorganisms associated with Deterioration of Tomato (*Lycopersicon esculentum*) and Pawpaw (*Carica papaya*) Fruits. *International Journal of Current Microbiology and Applied Science*. Vol 3 (5); 501-512.
- McGlynn, W. (2003). The Importance of Food pH in Commercial Canning Operation. *Oklahoma Cooperative Extension Service*.
- Moneruzzaman, K, M. Hossain, S, B, M, S. Sani. Saifuddin. (2008). Effect of Stages of Maturity and Ripening Conditions on the Biochemical Characteristics of Tomato. *American Journal of Biochemistry and Biotechnology*. Vol 4(4): 336-344.
- Novita, M. Satriana. Etria, H. (2015). Kandungan Likopen Dan Karotenoid Buah Tomat (*Lycopersicum Pyriforme*) Pada Berbagai Tingkat Kematangan: Pengaruh Pelapisan Dengan Kitosan Dan Penyimpanan. *Jurnal teknologi dan Industri Pertanian Indonesia*. Vol. 7 No.1.
- Nugraheni, S. Endrasari, R. Indrie, A. (2017). Perbedaan Kualitas Buah Tomat Hasil Produksi *Screen House* dan Lahan Terbuka Selama Penyimpanan. *Balai Pengkajian teknologi Pertanian Jawa Tengah*. Semarang.
- OECD. (2005). Guidance on Objective Tests to Determine Quality of Fruits and Vegetables and Dry and Dried Produce. *International Standardisation Of Fruit and Vegetables*.
- Pangloli P, Hung Y-C. (2011). Efficacy of slightly acidic *electrolyzed water* in killing or reducing *Escherichia coli* O157:H7 on Iceberg lettuce and tomatoes under simulated food service operation conditions. *J. Food Sci* Vol 76(6):M361-366.

- Park, H. Hung, Y. Donghwan, C. (2004). Effects of Chlorine and pH on Efficacy of Electrolyzed Water for Inactivating *Escherichia coli* O157:H7 and *Listeria monocytogenes*. *Internaional Journal of Food Microbioligy* 92:13-18.
- Peralta, I, E. Spooner, D.M. (2000). Classification of Wild Tomatoes : a riview. *Kurtizia* Vol 76(1):45-54.
- Peraturan Pemerintah Republik Indonesia. (2004). Keamanan, Mutu dan Gizi Pangan. Jakarta.
- Purwadi, A. Usada, W. Isyuniarto. (2007). Pengaruh Lama Waktu Ozonisasi Terhadap Umur Simpan Buah Tomat (*Lycopersicum esculentum* Mill). Pustek Akselerator dan Proses Bahan-BATAN.
- Rahman. SME., Imran. K., Deog-Hwan, Oh. (2016). Electrolyzed water as Novel Sanitizer in The Food Industry: Current Trends and Future Perspectives. *Comprehensive Review In Food Science and Food Safety*; 15.
- Saenmuang, S. Al-Haq, M, I. Yoshio, M. Yoshinori, K. Seiichi, O. (2012). Particle Size Distribution Of Nano-Mist In A Spinach-Storage Atmosphere And Its Effect On Respiration And Qualities. *Journal of Food Engineering* 1(12);69-77.
- Scientific Committee on Food. (2002). Risk People on The Microbiological Contamination of Fruits and Vegetables Eaten Raw. *European Commission : Health & Consumer Protection Directorate General*. Brussle, Belgium.
- Setiawan, D. Sibarani, J. Iryanti, E, S. (2013). Perbandingan Efektifitas Disinfektan Kaporit, Hidrogen Peroksida dan Pereaksi Feton ($\text{H}_2\text{O}_2/\text{Fe}^{2+}$). *Cakra Kimia*. Vol 1(2).
- Shin J. and Susan E. M. Selke. (2014). *Food Processing: Principles and Applications*, Second Edition Chapter 11: Food Packaging. John Wiley and Sons, Ltd.
- Sood, M. kaul, R, K. Smriq, S. (2011). Effect of Post Harvest Treatments on Changes in Sugar and Lycopene Content of Tomato (*Lycopersicon esculentum*). *World Journal of Agricultural Sciences*. Vol 7 (5): 613-616.
- Standar nasional Indonesia (SNI 3553:2015). (2015). Air Mineral. Badan Stadnarisasi Nasional. Jakarta.
- Sudarmadji, S. and Haryono, B. (1989). Prosedur Karakteristik Untuk Bahan Makanan dan Pertanian. *Liberty*. Yogyakarta
- Susenas. (2016). Konsumsi Buah dan Sayur Dalam Rangka Hari Gizi Nasional 2017. Jakarta.
- Susiwi. (2009). Handout Kerusakan Pangan. Jurusan Pendidikan Kimia. Universitas Pendidikan Indonesia.

- Suslow, T. (1997). Postharvest Chlorination : Basic Properties and Key Points for Effective Disinfection. Extension Specialist, Department of Vegetable Crops, University of California, Davis.
- Teka, A, T. (2013). Analysis of The Effect of Maturity Stage on The Postharvest Biochemical Quality Characteristic of Tomato (*Lycopersicon esculentum* Mill.) Fruit. Department of Postharvest Management.College of Agriculture and Veterinary Medicine. Jimma University.Jimma, Ethiopia.3(5);180-186.
- Thai Agricultural Standard*(TAS 1530-2007). (2007). Tomato. National Bureau of Agricultural Commodity and Food Standard. Ministry of Agriculture and Cooperatives.
- USDA. (2005). Tomatoes Shipping Point and Market Inspection Instructions. United States Department of Agriculture, Agricultural Marketing Service, Washington DC.
- USDA. (1991). United States Standard for grades of fresh tomatoes. United States Department of Agriculture, Agricultural Marketing Service, Washington DC.
- USDA National Nutrient database. (2016). Basic Report: 11529, Tomatoes, red, ripe, raw, year round average. National Agriculture Library.
- USDA. (2015). Hypochlorous Acid Handling. United States Department of Agriculture, Agricultural Marketing Service, Washington DC.
- Wisniewsky,A.M., A.G. Bonita, L.G. Mark, C.A. Reitmeier. (2000). Reduction of *Escherichia coli* O157:H7 counts on whole fresh apples by treatment with sanitizers. *Journal of Food Protection*, V 63, No. 6.
- Yahia, E, M. Xiuming, H. Athanasios, P, P. (2005). Infulence Of Crop Management Decision On Postharvest Quality of Greenhouse Tomatoes. Faculty of Chemistry, Autonomous of Queretaro. Mexico.
- Žnidarčič, D. Tomaž, P. (2006). Comparative Study of Quality Changes in Tomato cv. 'Malike' (*lycopersicum esculentum* Mill) whilst stored at different temperatures. *Acta Agriculture Slovenica*. 82-2;235-243.